

Total Ques: 45
Duration : 1 hr

NEET 2018

1

A

Agarwal Bandhu Gyan Kendra

12, Surya Nagar, Agra

NEET 2018 CHEMISTRY PAPER

Date: 27/07/2018
Test Booklet Code: PAP-0352
Test Booklet No: 521830

AGARWAL BANDEHU GYAN KENDRA AGARWAL BANDEHU GYAN KENDRA AGARWAL BANDEHU GYAN KENDRA AGARWAL BANDEHU GYAN KENDRA AGARWAL BANDEHU GYAN KENDRA

- The correct difference between first and second-order reactions is that :
 - the half-life of a first-order reaction does not depend on $[A]_0$; the half-life of a second-order reaction does depend on $[A]_0$
 - a first-order reaction can be catalyzed; a second-order reaction cannot be catalyzed
 - the rate of a first-order reaction does not depend on reactant concentrations; the rate of a second-order reaction does depend on reactant concentrations
 - the rate of a first-order reaction does depend on reactant concentrations; the rate of a second-order reaction does not depend on reactant concentrations
- Among CaH_2 , BeH_2 , BaH_2 , the order of ionic character is :
 - $\text{CaH}_2 < \text{BeH}_2 < \text{BaH}_2$
 - $\text{BeH}_2 < \text{BaH}_2 < \text{CaH}_2$
 - $\text{BeH}_2 < \text{CaH}_2 < \text{BaH}_2$
 - $\text{BaH}_2 < \text{BeH}_2 < \text{CaH}_2$
- In which case is the number of molecules of water maximum ?
 - 0.18 g of water
 - 0.00224 L of water vapours at 1 atm and 273 K
 - 18 mL of water
 - 10^{-3} mol of water
- Consider the change in oxidation state of Bromine corresponding to different emf values as shown in the diagram below :

$$\begin{array}{c}
 \text{BrO}_4^- \xrightarrow{1.82 \text{ V}} \text{BrO}_3^- \xrightarrow{1.5 \text{ V}} \text{HBrO} \\
 \text{Br}^- \xleftarrow{1.0652 \text{ V}} \text{Br}_2 \xleftarrow{1.595 \text{ V}}
 \end{array}$$

Then the species undergoing disproportionation is :

 - BrO_4^-
 - Br_2
 - BrO_3^-
 - HBrO
- In the structure of ClF_3 , the number of lone pairs of electrons on central atom 'Cl' is :
 - two
 - four
 - one
 - three
- The correct order of N-compounds in its decreasing order of oxidation states is :
 - $\text{HNO}_3, \text{NO}, \text{NH}_4\text{Cl}, \text{N}_2$
 - $\text{HNO}_3, \text{NH}_4\text{Cl}, \text{NO}, \text{N}_2$
 - $\text{HNO}_3, \text{NO}, \text{N}_2, \text{NH}_4\text{Cl}$
 - $\text{NH}_4\text{Cl}, \text{N}_2, \text{NO}, \text{HNO}_3$
- Which one of the following elements is unable to form MF_6^- ion ?
 - Al
 - B
 - Ga
 - In
- The correct order of atomic radii in group 13 elements is :
 - $\text{B} < \text{Al} < \text{Ga} < \text{In} < \text{Tl}$
 - $\text{B} < \text{Ga} < \text{Al} < \text{Tl} < \text{In}$
 - $\text{B} < \text{Al} < \text{In} < \text{Ga} < \text{Tl}$
 - $\text{B} < \text{Ga} < \text{Al} < \text{In} < \text{Tl}$
- Considering Ellingham diagram, which of the following metals can be used to reduce alumina ?
 - Zn
 - Mg
 - Fe
 - Cu
- Which of the following statements is not true for halogens ?
 - All are oxidizing agents
 - All but fluorine show positive oxidation states
 - All form monobasic oxyacids
 - Chlorine has the highest electron-gain enthalpy
- Regarding cross-linked or network polymers, which of the following statements is *incorrect* ?
 - They are formed from bi- and tri-functional monomers
 - Examples are bakelite and melamine
 - They contain covalent bonds between various linear polymer chains
 - They contain strong covalent bonds in their polymer chains
- The difference between amylose and amylopectin is :
 - Amylose have $1 \rightarrow 4$ α -linkage and $1 \rightarrow 6$ β -linkage
 - Amylopectin have $1 \rightarrow 4$ α -linkage and $1 \rightarrow 6$ β -linkage
 - Amylopectin have $1 \rightarrow 4$ α -linkage and $1 \rightarrow 6$ α -linkage
 - Amylose is made up of glucose and galactose

- 13 Nitration of aniline in strong acidic medium also gives m-nitroaniline because :
- In electrophilic substitution reactions amino group is meta directive
 - In absence of substituents nitro group always goes to m-position
 - In spite of substituents nitro group always goes to only m-position
 - In acidic (strong) medium aniline is present as anilinium ion

- 14 Which of the following oxides is most acidic in nature ?
- BeO
 - BaO
 - MgO
 - CaO


- 15 A mixture of 2.3 g formic acid and 4.5 g oxalic acid is treated with conc. H_2SO_4 . The evolved gaseous mixture is passed through KOH pellets. Weight (in g) of the remaining product at STP will be :
- 3.0
 - 2.8
 - 1.4
 - 4.4

- 16 The compound A on treatment with Na gives B, and with PCl_5 gives C, B and C react together to give diethyl ether. A, B and C are in the order :
- $C_2H_5OH, C_2H_5Cl, C_2H_5ONa$
 - $C_2H_5Cl, C_2H_6, C_2H_5OH$
 - $C_2H_5OH, C_2H_6, C_2H_5Cl$
 - $C_2H_5OH, C_2H_5ONa, C_2H_5Cl$

- 17 Hydrocarbon (A) reacts with bromine by substitution to form an alkyl bromide which by Wurtz reaction is converted to gaseous hydrocarbon containing less than four carbon atoms. (A) is :
- $CH_2 = CH_2$
 - $CH_3 - CH_3$
 - $CH \equiv CH$
 - CH_4

- 18 Which oxide of nitrogen is **not** a common pollutant introduced into the atmosphere both due to natural and human activity ?
- NO_2
 - N_2O
 - N_2O_5
 - NO


- 19 The compound C_7H_8 undergoes the following reactions :




- The product 'C' is :
- o*-bromotoluene
 - 3-bromo-2, 4, 6-trichlorotoluene
 - m*-bromotoluene
 - p*-bromotoluene

- 20 Which of the following molecules represents the order of hybridisation sp^2, sp^2, sp, sp from left to right atoms ?
- $CH_2 = CH - C \equiv CH$
 - $CH_2 = CH - CH = CH_2$
 - $HC \equiv C - C \equiv CH$
 - $CH_3 - CH = CH - CH_3$

- 21 Which of the following carbocations is expected to be most stable ?



- 22 Which of the following is correct with respect to -I effect of the substituents ? (R = alkyl)
- $-NR_2 < -OR < -F$
 - $-NH_2 > -OR > -F$
 - $-NH_2 < -OR < -F$
 - $-NR_2 > -OR > -F$







- the electrophile involved is :
- Formyl cation (CHO^+)
 - dichloromethyl anion ($CHCl_2^-$)
 - dichloromethyl cation ($CHCl_2^+$)
 - dichlorocarbene ($:CCl_2$)


- 24 Carboxylic acids have higher boiling points than aldehydes, ketones and even alcohols of comparable molecular mass. It is due to their :
- formation of carboxylate ion
 - more extensive association of carboxylic acid via van der Waals force of attraction
 - formation of intramolecular H-bonding
 - formation of intermolecular H-bonding










5
2
1
8
3
0

25 Compound A, $C_9H_{10}O$, is found to react with NaOI (produced by reacting Y with NaOH) and yields a yellow precipitate with characteristic smell. A and Y are respectively

- (a)  and I_2
- (b)  and I_2
- (c)  and I_2
- (d)  and I_2

26 Identify the major products P, Q and R in the following sequence of reactions :



- | | | |
|---|---|--|
| P | Q | R |
| (a)  |  |  |
| (b)  |  | $\text{CH}_3\text{CH}(\text{OH})\text{CH}_3$ |
| (c)  |  | $\text{CH}_3\text{CH}_2\text{-OH}$ |
| (d)  |  | $\text{CH}_3\text{-CO-CH}_3$ |


27 Which of the following compounds can form a zwitterion ?

- (a) Acetanilide
- (b) Benzoic acid
- (c) Aniline
- (d) Glycine


28 The correction factor 'a' to the ideal gas equation corresponds to :

- (a) volume of the gas molecules
- (b) electric field present between the gas molecules
- (c) density of the gas molecules
- (d) forces of attraction between the gas molecules

29 For the redox reaction



the correct coefficients of the reactants for the balanced equation are :



- (a) 2 5 16
- (b) 2 16 5
- (c) 16 5 2
- (d) 5 16 2


30 When initial concentration of the reactant is doubled, the half-life period of a zero order reaction :

- (a) is doubled
- (b) is tripled
- (c) is halved
- (d) remains unchanged

31 The bond dissociation energies of X_2 , Y_2 and XY are in the ratio of 1 : 0.5 : 1. ΔH for the formation of XY is -200 kJ mol^{-1} . The bond dissociation energy of X_2 will be :

- (a) 100 kJ mol^{-1}
- (b) 800 kJ mol^{-1}
- (c) 200 kJ mol^{-1}
- (d) 400 kJ mol^{-1}

32 Which one of the following conditions will favour maximum formation of the product in the reaction :




- (a) Low temperature and low pressure
- (b) High temperature and high pressure
- (c) Low temperature and high pressure
- (d) High temperature and low pressure

33 Iron exhibits bcc structure at room temperature. Above 900°C , it transforms to fcc structure. The ratio of density of iron at room temperature to that at 900°C (assuming molar mass and atomic radii of iron remains constant with temperature) is :

- (a) $\frac{4\sqrt{3}}{3\sqrt{2}}$
- (b) $\frac{3\sqrt{3}}{4\sqrt{2}}$
- (c) $\frac{\sqrt{3}}{\sqrt{2}}$
- (d) $\frac{1}{2}$

34 Consider the following species :



Which one of these will have the highest bond order ?

- (a) CN^-
- (b) CN^+
- (c) NO
- (d) CN


5
2
1
8
3
0

35 Magnesium reacts with an element (X) to form an ionic compound. If the ground state electronic configuration of (X) is $1s^2 2s^2 2p^3$, the simplest formula for this compound is :

- (a) MgX_2 (b) Mg_2X
(c) Mg_2X_3 (d) Mg_3X_2





36 Which one is a wrong statement ?

- (a) An orbital is designated by three quantum numbers while an electron in an atom is designated by four quantum numbers
(b) The electronic configuration of N atom is



- (c) Total orbital angular momentum of electron in 's' orbital is equal to zero
(d) The value of m for d_{z^2} is zero

37 Following solutions were prepared by mixing different volumes of NaOH and HCl of different concentrations :



pH of which one of them will be equal to 1 ?

- (a) 1
(b) 4
(c) 2
(d) 3

38 On which of the following properties does the coagulating power of an ion depend ?

- (a) Size of the ion alone
(b) Both magnitude and sign of the charge on the ion
(c) The magnitude of the charge on the ion alone
(d) The sign of charge on the ion alone

39 Given van der Waals constant for NH_3 , H_2 , O_2 and CO_2 are respectively 4.17, 0.244, 1.36 and 3.59, which one of the following gases is most easily liquefied ?

- (a) H_2 (b) O_2
(c) NH_3 (d) CO_2

40 The solubility of $BaSO_4$ in water is $2.42 \times 10^{-3} \text{ g L}^{-1}$ at 298 K. The value of its solubility product (K_{sp}) will be :

(Given molar mass of $BaSO_4 = 233 \text{ g mol}^{-1}$)

- (a) $1.08 \times 10^{-12} \text{ mol}^2 \text{ L}^{-2}$
(b) $1.08 \times 10^{-14} \text{ mol}^2 \text{ L}^{-2}$
(c) $1.08 \times 10^{-10} \text{ mol}^2 \text{ L}^{-2}$
(d) $1.08 \times 10^{-8} \text{ mol}^2 \text{ L}^{-2}$

41 The type of isomerism shown by the complex $[CoCl_2(en)_2]$ is :

- (a) Coordination isomerism
(b) Ionization isomerism
(c) Geometrical isomerism
(d) Linkage isomerism

42 Which one of the following ions exhibits d-d transition and paramagnetism as well ?

- (a) $Cr_2O_7^{2-}$ (b) MnO_4^-
(c) CrO_4^{2-} (d) MnO_4^{2-}

43 Iron carbonyl, $Fe(CO)_5$ is :

- (a) mononuclear (b) trinuclear
(c) tetranuclear (d) dinuclear

44 Match the metal ions given in Column-I with the spin magnetic moments of the ions given in Column-II and assign the **correct** code :

Column-I	Column-II
(1) Co^{3+}	(i) $\sqrt{8}$ B.M.
(2) Cr^{3+}	(ii) $\sqrt{35}$ B.M.
(3) Fe^{3+}	(iii) $\sqrt{3}$ B.M.
(4) Ni^{2+}	(iv) $\sqrt{24}$ B.M.
	(v) $\sqrt{15}$ B.M.

(1)	(2)	(3)	(4)
(a) (i)	(ii)	(iii)	(iv)
(b) (iv)	(i)	(ii)	(iii)
(c) (iv)	(v)	(ii)	(i)
(d) (iii)	(v)	(i)	(ii)

45 The geometry and magnetic behaviour of the complex $[Ni(CO)_4]$ are :

- (a) tetrahedral geometry and diamagnetic
(b) square planar geometry and paramagnetic
(c) square planar geometry and diamagnetic
(d) tetrahedral geometry and paramagnetic

A										
Test Booklet No.			Date			Paper Code			Paper Description	
521830			27 - Jul - 18			PAP-0352			NEET 2018	
(1) a	(2) c	(3) c	(4) d	(5) a	(6) c	(7) b	(8) d	(9) b	(10) c	
(11) d	(12) c	(13) d	(14) a	(15) b	(16) d	(17) d	(18) c	(19) c	(20) a	
(21) b	(22) c	(23) d	(24) d	(25) b	(26) d	(27) d	(28) d	(29) a	(30) a	
(31) b	(32) c	(33) b	(34) a	(35) d	(36) b	(37) d	(38) b	(39) c	(40) c	
(41) c	(42) d	(43) a	(44) c	(45) a						